

NUMERICAL STUDY OF PLASMON PROPERTIES IN THE SU(2)-HIGGS MODEL

Jan SMIT

*Institute of Theoretical Physics, University of Amsterdam,
Valckenierstraat 65, 1018 XE Amsterdam, the Netherlands*

We discuss an explorative computation of real time autocorrelation functions, in the classical approximation. The results for the ‘plasmon’ frequencies and damping rates appear compatible with the divergencies expected from perturbation theory.

1 Motivation

The (effectively) classical approximation has been used in computations of the Chern-Simons diffusion rate in the SU(2)-Higgs model, which is an essential ingredient in theories of baryogenesis. It is potentially a very useful approximation for non-equilibrium processes in field theory. However, its formal properties are not very well known yet and it is not clear either how well the classical field theory approximates the quantum theory at high temperatures.

These issues can be studied in the weak coupling expansion. The classical theory has divergencies at finite temperature and the question naturally arises if the theory is renormalizable in some sense. The answer is affirmative for ϕ^4 theory and the hot classical theory can approximate the quantum theory well at weak coupling¹. For gauge theories the situation is more complicated. Their leading hard thermal loop effects are non-local and counterterms would have to have this property as well (although the specific form of the nonlocalities may be representable by locally coupled new degrees of freedom).

Still, one can suppose that any divergent quantity will be unobservable at low energies. Divergent frequencies and damping rates will be unobservable at sufficiently large time scales. The question is then, if there remain physically interesting phenomena at such time scales.

For time independent quantities the classical approximation is a form of the dimensional reduction approximation. However, for the time component A_0 of the gauge field the coefficients of $\text{Tr } A_0^2$ and $\text{Tr } A_0^4$ are zero in the classical theory², while they have well defined values in the dimensionally reduced theory, including a Debye mass counterterm $\propto \text{Tr } A_0^2$. The absence of this counterterm in the classical theory implies that the corresponding Debye mass is divergent. For static quantities this is not a problem since A_0 simply decouples. Such a decoupling approximation is often made in addition to dimensional reduction because the perturbative Debye mass is relatively large. For dynam-

ical properties one may hope that a similar decoupling is possible in the sense describe above.

The situation is more serious in the dynamical case since also the classical plasmon frequency associated with the modes A_k , $k = 1, 2, 3$, is divergent³. This can be seen in a one loop calculation on the lattice. The two phenomena (diverging Debye mass and plasmon frequency) are related by the fact that there is no local counterterm available for A_μ in the classical lagrangian.

So it seems a good idea to compute nonperturbatively various correlation functions in the classical model and study their lattice spacing dependence. The results can then also be compared with analytic calculations, in the classical and quantum theory. This work was undertaken with Wai Hung Tang⁴. We concentrated on autocorrelation functions

$$C_O(t) = \langle O(t)O(0) \rangle \quad (1)$$

of gauge invariant observables with the quantum numbers of the Higgs and W fields, at zero momentum,

$$O \rightarrow H = \varphi^\dagger \varphi, \quad \rightarrow W_k^\alpha = i \text{Tr} \phi^\dagger (\partial_k - i A_k^\beta t_\beta) \phi t_\alpha, \quad (2)$$

with φ the Higgs doublet, ϕ its matrix version ($i\tau_2 \varphi^*, \varphi$) and $t_\alpha = \tau_\alpha/2$ the $SU(2)$ generators. In the Higgs phase these fields correspond to the usual Higgs and gauge fields. Since $\phi = \sqrt{H} V$ with $V \in SU(2)$, W_μ^α is essentially the gauge field in the unitary gauge $V = 1$, as can be seen in perturbation theory by replacing φ by its vacuum expectation value, neglecting fluctuations. In the plasma phase this reasoning breaks down as the theory behaves like QCD and perturbation theory runs into infrared problems.

2 Simulation

The effective classical hamiltonian on a spatial lattice can be given schematically in the form²

$$\begin{aligned} \frac{H_{\text{eff}}}{T} = & \bar{\beta} \sum_{\mathbf{x}} \left[\frac{1}{2} z_E \bar{E}_{m\mathbf{x}}^\alpha \bar{E}_{m\mathbf{x}}^\alpha + z_\pi \bar{\pi}_{\mathbf{x}}^\dagger \bar{\pi}_{\mathbf{x}} + \sum_{m < n} \left(1 - \frac{1}{2} \text{Tr} U_{mn\mathbf{x}} \right) \right. \\ & \left. + (D_m \bar{\varphi}_{\mathbf{x}})^\dagger D_m \bar{\varphi}_{\mathbf{x}} + \bar{\lambda} (\bar{\varphi}_{\mathbf{x}}^\dagger \bar{\varphi}_{\mathbf{x}} - \bar{v}^2)^2 \right]. \end{aligned} \quad (3)$$

We use the parametrization and numerical implementation of Krasnitz⁵. Comparison with dimensional reduction leads to the following identification of parameters²

$$\bar{\beta} \approx \frac{4}{g^2 a T}, \quad \bar{\lambda} \approx \frac{m_H^2}{2m_W^2}, \quad (4)$$

$$4\bar{\lambda}\bar{v}^2 = a^2 T^2 \left[\frac{m_H^2}{T^2} - \frac{9g^2}{4} \left(\frac{3+\rho}{18} - \frac{3+\rho}{3} \frac{2\Sigma}{aT} \right) - \frac{81g^4}{16} \right. \\ \left. \times \frac{1}{8\pi^2} \left(\frac{149+9\rho}{486} + \frac{27+6\rho-\rho^2}{27} \ln \frac{aT}{2} - \frac{27\eta+6\rho\bar{\eta}-\rho^2\tilde{\eta}}{27} \right) \right], \quad (5)$$

where a is the lattice spacing, η , $\bar{\eta}$ and Σ are numerical constants, $\rho = m_H^2/m_W^2$, and g^2 , $m_{H,W}^2$, are running couplings and masses in the 4D quantum theory in the $\overline{\text{MS}}$ scheme with scale parameter $\mu = 4\pi e^{-\gamma}T \approx 7T$. Furthermore, $z_E/z_\pi = 1$ to a good approximation, which leaves only one parameter $z = z_E = z_\pi$ to set the time scale relative to the lattice spacing. Intuitively one expects $z = 1 + O(g^2, \lambda)$, but recent work suggests that z may have to be taken substantially different from 1, depending on the lattice spacing³.

We used $z = 1$, $g^2 = 4/9$, $\rho = 2\bar{\lambda} = 1$, and various $\bar{\beta}$, $\bar{v}^2$ and lattice sizes N^3 . First we did a scan with $N = 24$ and $\bar{v}^2$ fixed, implying varying T/m_H by eq. (5). The ratio T/m_H can be converted to T/T_c since the transition point T_c/m_H is known reasonably well. Then we spent more effort in the plasma phase with $(N, \bar{\beta}) = (20, 11)$ and $(32, 18.3)$, while $\bar{v}^2$ was adjusted such that $T/T_c = 1.52$ in both cases. Hence, the lattice spacing changed by a factor of $11/18.3 = 0.60$ while the physics remained the same according to (5), including the physical volume: $LT = NaT = 9N/\bar{\beta} \approx \text{const}$. In the Higgs phase similar effort was spent with $(N, \bar{\beta}) = (20, 13)$ and $(32, 21.7)$, without adjusting $\bar{v}^2$, however, which led to somewhat different T/T_c : 0.85 and 0.78, respectively.

Fig. 1 shows examples of correlation functions. On the left we see $C_H(t)$ in the Higgs phase for $\bar{\beta} = 13$ and 21.7. The damped oscillations can be fitted to the pole dominance form

$$Re^{-\gamma t} \cos(\omega t + \alpha) + \text{const}, \quad (6)$$

with $\alpha, \text{const} \approx 0$. The plot for $C_W(t)$ looks similar, but with more damping and the additive constant effectively nonzero. In the plasma phase the results are qualitatively different. There $C_H(t)$ still shows oscillations with physical ($a\omega < 1$) frequencies, but with a much larger effective damping coefficient γ , and the additive background drops in time. For $C_W(t)$ we find only a few oscillations at unphysically short times (t/a of order 2), after which a slow roughly exponential decay sets in ($\omega \rightarrow 0$ and small γ in (6)), shown on the right in Fig. 1. Fig. 2 shows the results for ω and γ in units of g^2T (in lattice units, all the $a\omega$ and $a\gamma$ are reasonable small compared to 1).

3 Interpretation

Consider first the frequencies $\omega_{H,W}$ in the Higgs phase, where H and W correspond to the usual Higgs and W fields in leading order perturbation theory. At




Figure 1: Left: $C_H(t)$ in the Higgs phase for $\bar{\beta} = 13$ (small amplitude) and 21.7 (large amplitude). Right: plot of $\ln C_W(t)$ in the plasma phase. The crossing lines correspond to $\bar{\beta} = 11, 18.3$ at $T/T_c = 1.52$. The top curve corresponds to $\bar{\beta} = 11$ at a different $\bar{v}^2$ such that $T/T_c = 1.23$. The time is in lattice units: ‘time’ = t/a . The insert shows $\ln C_w(t)$ versus $\ln t/a$.

first sight the small deviations of the 32^3 data from the other data (interpolated by eye) suggest lattice spacing independence. This is what we optimistically highlighted in the first version of the paper⁴. However, our understanding increased substantially since then and as a result (being pressed as well by the referees of our paper) we reanalyzed the data, in the spirit of the expected $1/\sqrt{a}$ divergence in ω_W from one loop perturbation theory. The upshot (expressed in the revised version) is that the data are in fact compatible with this divergence. A recent analysis shows that also their magnitude can be understood from perturbation theory⁶.

The damping rates on the other hand are expected to be finite to leading nontrivial order. In the Higgs phase they compare well with the analytic results in the quantum theory. For example, the $\bar{\beta} = 21.7$ ($T/T_c = 0.78$) data give $\gamma_W/g^2T = 0.22(4)$ and $\gamma_H/g^2T = 0.028(4)$, to be compared with 0.176 and ≈ 0.02 , respectively⁷. In the plasma phase there are strong fluctuations in $C_H(t)$ causing large errors on γ_H ; the errors on γ_W are reasonable. In both phases the damping rate data are compatible with lattice spacing independence, which, given the large errors is not a very strong statement. However, the a independence of γ_W in the plasma phase ($\gamma_W/g^2T = 0.31(2), 0.028(4)$ for respectively $\bar{\beta} = 11, 18.3$) suggests that it is a physical quantity as well.




Figure 2: ‘Plasmon’ frequencies ω/g^2T (left) and damping rates γ/g^2T (right) versus T/T_c . Solid symbols correspond to H , open symbols to W ; circle: 24^3 data; diamond: 20^3 data; square: 32^3 data.

A more complete understanding may be expected from analysis of the full autocorrelation functions in perturbation theory. Work in this direction is in progress⁸. For example, the qualitative difference between $C_H(t)$ and $C_W(t)$ at short times can be explained this way⁶. A good understanding of the (over-?) damping in $C_W(t)$ at larger times may also be obtainable this way.

Acknowledgements

We thank Gert Aarts and Alex Krasnitz for useful conversations. This work is supported by FOM and NCF/NWO.

References

1. G. Aarts and J. Smit, *Phys. Lett. B* **393**, 395 (1997); G. Aarts and J. Smit, hep-ph/9707342; G. Aarts, these proceedings.
2. W.H. Tang and J. Smit, *Nucl. Phys. B* **482**, 265 (1996).
3. P. Arnold, *Phys. Rev. D* **55**, 7781 (1997).
4. W.H. Tang and J. Smit, hep-lat/9702017.
5. A. Krasnitz, *Nucl. Phys. B* **455**, 320 (1995).
6. B. Bökeler and M. Laine, hep-ph/9707489.
7. E. Braaten and R.D. Pisarski, *Phys. Rev. D* **42**, 2156 (1990); T.S. Biró and M.H. Thoma, *Phys. Rev. D* **54**, 3465 (1996).
8. P. Arnold, private communication.